

NOTES AND ERRATA TO JOURNAL OF SOUTH AFRICAN BOTANY, SUPPLEMENTARY VOLUME I (1944); THE GENUS *OXALIS* IN SOUTH AFRICA

BY CAPTAIN T. M. SALTER, R.N.

Page

- 25 Penultimate line, for 266 read 226.
- 31 Insert after par. 1: For notes on the formation of the bulbs and the action of the contractile root in seedlings of *Oxalis* producing endospermous seeds, see *Journal of South African Botany*, Vol. XVII (1951), p. 190.
- 31 Par. 2, last 4 lines. See Notes on *O. laburnifolia* Jacq., Part II, p. 18, par. 3. (*Journ. of S.A. Bot.* XIV (1948).)
- 35 Par. 3, last 2 lines. See Notes on *O. laburnifolia* Jacq., Part II, p. 15, Aerial bulbils.
- 38 Line 5, for *Sessilifoliae* read *Sessilifoliatae*.
- 55 In key to Group 2, for (101) *fabaeifolia* read (111).
- 64 In key to Group 10, after line 3 insert:

Petioles flattened: plant producing aerial bulbils . . . (56a) *laburnifolia*
 Petioles not flattened: plant not producing aerial
 bulbils (56) *purpurea*
- 75 Line 3, for cuneate-obovate read cuneate-obcordate.
- 75 Var. **purpurascens** Salter, par. 2, line 2, for petioles read pedicels.
- 87 Penultimate line, for S.A.M. 49572 read 49752.
- 124 **O. luteola** var. **minor** Salter, line 2, for 1·5—4 cm. read mm.
- 129 Par. 2, last line, before near Woodville, insert GEORGE DIV.
- 131 Line 4, for *S* 4789 read 4879.
- 140 Omit **O. purpurea** L. Forms (i), (ii) and (iii). See Notes on *O. laburnifolia* Jacq. and its variety *latifolia*. (*Journ. of S.A. Bot.* XIV (1948), Part I.)
- 142 After Species 56 insert Species 56a:

56a. **O. laburnifolia** Jacq. *Oxal.* (1794) 63, tab. 28—*O. laburnifolia* var. *angustata* Sond.—*O. sanguinea* Jacq., l.c. 64, tab. 29—*O. arthropphylla* Turcz.—*O. purpurea* L. Form E (iii) Salter. Stemless, 3—6 cm. high. *Bulb* broadly or narrowly ovoid, 2—4 cm. long, acute at the apex, the tunics smooth, dark brown: *aerial bulbils* small, very numerous sometimes several hundreds, densely massed in the leaf axils, fully developed about two months after flowering. *Rhizome* usually short. *Leaves* 4—10, spreading:

petioles stout, hispid, somewhat flattened, channelled on the upper side, often red-tinged: leaflets 3, 2—4 cm. long, sparsely pubescent, ciliate, more or less blood-red beneath, dotted with numerous small black streaks when dried, obtuse, the medial oblong-oblancoelate, the lateral narrower, oblique at the base. *Peduncles* 1-fl., hirsute, shorter than the petioles, with 2 alternate bracts below the middle. *Sepals* lanceolate, 4—5 mm. long, with longitudinal black streaks when dried. *Corolla* 2·5—3·5 cm. long, yellow, with a narrowly cyathiform concolorous tube: lamina of the petals cuneate-obovate, attenuate to a rather short claw. *Filaments* glandular-pilose, the longer 3·5—6 mm. long, dentate. *Ovary* callose, pubescent on the upper half, the chambers many-ovuled. *Seeds* endospermous.

VAN RHYNSDORP DIV.; between Olifants River and Knackisberg, *Zey.* 225; Ouplaas Hills, west of Klaver, *Salter* 5445; 24 miles north of Lamberts Bay, *Nortier* (Kirstenbosch 263/46 cult.); *Leipoldt* 4296. Fl. May—June.

Var. *latifolia* Salter, Forms A and B are described in Journ. of S.A. Bot., Vol. XIV (1948), pp. 14, 15.

- 175 Penultimate line, for *Sessilifoliae* read *Sessilifoliatae*.
 191 *O. stenoptera* var. *undulata*, par. 4, line 3, for corolla read calyx.
 238 *O. hirta* Form A, line 1, for tab. 53 read tab. 13.
 239 *O. hirta* Form B, par. 2, line 3, for *Phillips* 9495 read 9475.
 241 Lines 22, 23, for var. *intermedia* read var. *polioeides*.
 266 Par. 2, line 1, after Louwpoort for *S* 5489 read 4589.
 267 Par. 1, line 3, after Plate VII insert Opp. p. 226.
 267 Var. *purpurea*, line 3 and var. *lilacea*, line 3, after Plate VIII insert Opp. p. 227.
 267 Par. 2, line 4, after CLANW. DIV.; for *S* 7346 read 7246.
 283 Par. 3, line 2, for S.A.M. 48376 read 48375.
 321 *O. variifolia*, line 2, for *stendodactyla* read *stenodactyla*.
 328 *O. ligulata*, in final note, for but it is represented, read and it is not represented. None of the poor specimens of Drège 3198 seen by me is the species described by Sonder, they appear to be probably Form B of *O. obtusa* Jacq.
 333 For 525 *commutata* Sond. read 526.
 334 For 658 *viscosa* E. Mey. ex Sond. read 668.
 349 For 8158 *hirta* var. *polioeides* read 8157.